


Chapter 15: Acid-Base Regulation

Defending Proton Homeostasis, Instructor Guide

Mechanisms and Logic in Human Physiology | Silldorff & Robinson | Kendall Hunt

Format: Left column = instructor narrative in the author's voice. Right column = structured outline with causal chains, key distinctions, common errors (red), and prediction prompts (gold italic). The organizing spine of this integration chapter: one parameter, body-fluid $[H^+]$, is defended by three layers (chemical buffers, the lungs, the kidneys) that all act on a single reaction, the carbonic acid equation, with the lungs steering its CO_2 end and the kidneys its bicarbonate end.



Companion to **Mechanisms and Logic in Human Physiology** by Erik P. Silldorff and Gerald D. Robinson (Kendall Hunt Publishing). The half-year and full-year web subscriptions and the print version are available at <https://he.kendallhunt.com/product/mechanisms-and-logic-human-physiology-2>. For those teaching A&P more classically in a 2-semester sequence, the book is also packaged with Visible Body to support the anatomical portions of those courses (365-day subscriptions): <https://he.kendallhunt.com/product/mechanisms-and-logic-human-anatomy-and-physiology-visible-body-1>.

Cite as: Silldorff, E. P., & Robinson, G. D. (2026). Chapter 15 – Acid-Base Regulation: An Instructor Guide for *Mechanisms and Logic in Human Physiology*. <https://doi.org/10.5281/zenodo.22834156>

This work is licensed under CC BY-NC-ND 4.0. <https://creativecommons.org/licenses/by-nc-nd/4.0/>

© 2026 Erik P. Silldorff and Gerald D. Robinson



[Resource Hub](#)

Section 1: The Defended Parameter — Why Protons Get the Tightest Setpoint in the Body

Continuation of Unit 4. This is an integration chapter: the controlled parameter is body-fluid $[H^+]$, and the controllers are the chemical buffers, the lungs, and the kidneys working as one system.

Instructor Narrative	Structured Outline / Causal Chain
The move: <i>The chapter is organized around the defense of a single parameter, body-fluid $[H^+]$. It requires this much regulatory machinery because protons alter the charge on protein side groups and destabilize</i>	Why This Chapter Sits in the Integration Unit Controlled parameter: body-fluid $[H^+]$, expressed as pH.

protein structure, and three lines of defense, chemical buffers, the lungs, and the kidneys, all act on that one parameter.

H⁺ is shorthand, and proteins are why it matters

Bare protons do not float free in solution. Instead, their extremely high positive charge density makes them instantly associate with water molecules, forming hydronium (H₃O⁺). H⁺, then, is shorthand for the more complex combination of protons and the water molecules they interact with. In pure water at 25 °C, only about 1 in 500 million water molecules dissociate to release a proton. In blood, the concentration of these ions is even lower due to the specific balance of acids and bases. What students need is not that chemistry but its consequence. Despite the very low proton concentration in body fluids, even slight deviations in their abundance can be deleterious. Protons alter the charge on amino acid side groups, and those altered charges change how a protein folds. Many proteins are only marginally stable, so a small shift in [H⁺] can unfold an enzyme, channel, or pump and kill the cell quickly. The parameter is abstract; the stakes are entirely concrete.

Regulated a million-fold tighter than sodium

Body-fluid proton concentration is held stable roughly a million times more closely than is Na⁺. Normal arterial pH (details to follow) sits in a window of about 7.35 to 7.45, a sliver of the 0-to-14 scale, and a full unit deviation either way is lethal. A typical Western diet of meats, grains, and dairy is a net acid producer, so this is not a system at rest: acid is made continuously in metabolism and must be matched, over time, by acid excretion.

Three defenses, one parameter: the spine of the chapter

The body defends pH in three layers that differ mainly in speed and finality: chemical buffers act instantly but only hide protons; the lungs adjust CO₂ within seconds but only blow off volatile acid (acids that can be transitioned to a gas, CO₂); the kidneys act over hours but are the only system that truly eliminates fixed acid (cannot be turned into a gas, e.g., lactic acid) and rebuilds buffer. Every mechanism in this chapter is one of those three layers acting on one key correlation, the carbonic acid equation (see below).

- Controllers: three organ-level systems acting together: chemical buffers, respiratory, renal.
- No single system fully solves the problem; integration is the point.

Why Protons Are So Dangerous

- Protons alter the charge on amino acid side groups, altering protein folding and function.
- Marginally stable proteins fail fast; cell death follows.
- Held ~ 1,000,000× tighter than Na⁺; normal pH 7.35–7.45.

Causal Chain: The Three Lines of Defense (Fast to Final)

Chemical buffers bind added protons within milliseconds, converting free H⁺ into buffer-bound H⁺ → free [H⁺] falls and the pH change is blunted, but the protons remain in the body and buffer capacity is consumed → the resulting fall in pH is sensed by chemoreceptors, ventilation rises, and CO₂ is exhaled within seconds, clearing the volatile acid (the fraction convertible to CO₂ gas). Fixed acids cannot be converted to a gas and exhaled, so they remain in the body → the kidneys secrete H⁺ into acidic urine and synthesize new HCO₃⁻ over hours, removing the fixed acid and restoring the consumed buffer

Common Errors

- **Error:** Thinking buffers fix the problem. They only mask added protons; the acid stays in the body until the kidney excretes it.
- **Error:** Assuming pH is loosely controlled because the numbers look tiny. The opposite is true: this is the most tightly defended ion and the pH scale is logarithmic (a relatively large change in [H⁺] can result in a very small alteration in pH).

Prediction Prompt

- **Predict:** A diet shifts from meat-heavy to vegetable-heavy. Predict the direction of the net metabolic acid load and what the kidney must do differently.

Section 2: Acids, Bases, and the Meaning of Strength

Instructor Narrative	Structured Outline / Causal Chain
The move: The Brønsted-Lowry definitions take one line: an acid donates a proton, a base accepts one. The term that carries the physiology is strength. Abundance is read off a concentration; strength is the tendency to release a proton, and it is what makes buffering possible. Donors, acceptors, and the conjugate pair An acid is a proton donor, a base a proton acceptor: $\text{HA} \leftrightarrow \text{H}^+ + \text{A}^-$, where the anion A^- left is the conjugate base. That is the whole vocabulary. The useful move is to read every acid as a pair, the protonated form and its conjugate base, because buffering is nothing more than a pair trading a proton back and forth. Strength is a tendency, and K_a measures it Two things set the $[\text{H}^+]$ a solution carries: how much acid is present (concentration) and how readily it dissociates (strength). Strength is captured by the acid dissociation constant, $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$. A strong acid like HCl dissociates almost completely: its K_a is enormous and its conjugate base (Cl^-) is so weak it binds essentially no protons. A weak acid like carbonic acid barely dissociates; its K_a is more than twelve orders of magnitude smaller than that of HCl. Physiology lives among the weak acids A buffer converts a strong, fully dissociating acid into a weak, barely dissociating one, trading a large pH change for a small one. Hold that	The Core Relationship • Acid = proton donor; base = proton acceptor: $\text{HA} \leftrightarrow \text{H}^+ + \text{A}^-$. • A^- = conjugate base; a strong acid has a weak conjugate base, and vice versa. • $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$; large K_a = strong acid. The Strength Spectrum (chapter examples) • HCl: very strong acid, yielding Cl^-, a very weak base (K_a very high). • H_2CO_3: weak acid, yielding HCO_3^-, a stronger base (K_a very low). • Water: very weak acid, yielding OH^-, a very strong base. Common Errors • Error: Conflating abundance with strength. A lot of a weak acid and a little of a strong acid can carry the same H^+ burden. • Error: Forgetting that a weak acid has a strong conjugate base (and vice versa): this inverse relationship drives buffering. Prediction Prompt

idea: it returns in the bicarbonate system, where infused strong acid is converted into weaker carbonic acid.

- **Predict:** Two 0.1 M acids are compared; one has the larger K_a . Predict which solution has the lower pH and which has the stronger conjugate base.

Section 3: The pH Scale — Why a Logarithm, and Why 0.1 Units Is a Big Deal

Instructor Narrative	Structured Outline / Causal Chain
The move: pH is a logarithm of proton concentration, and the logarithm is the source of most student misconceptions about pH. A shift from pH 7.35 to 7.45 is a substantial change in proton concentration rather than just a rounding difference, and the physiological significance of the rest of the chapter depends on that point. A logarithm tames an enormous range Proton concentrations span fourteen orders of magnitude and writing them in molarity is unwieldy. Sørensen's $\text{pH} = -\log[\text{H}^+]$ compresses that range into a 0-to-14 scale, and because body-fluid $[\text{H}^+]$ is always a small fraction of a molar, the negative log is always positive. pH 7 is neutral; below 7 is acidic (more H^+); above 7 is alkaline (less H^+). Small pH numbers correspond to large concentration changes Because the scale is logarithmic, a change of 0.1 pH unit is not small: moving from 7.45 to 7.35 is roughly a 26% increase in $[\text{H}^+]$, and a full unit change is a tenfold increase or decrease. The narrow clinical window of pH 7.35 to 7.45 therefore represents real, continuously compensated variation, not rounding noise. This band of blood pH compatible with life is just a tiny slice of the full scale.	The Definition and Its Logic • $\text{pH} = -\log[\text{H}^+]$; Sørensen, 1909; unit-less, ranges ~ 0–14. • Log chosen to compress 14 orders of magnitude into one sequence. • Higher $\text{H}^+ \rightarrow$ lower pH; pH 7 neutral, < 7 acidic, > 7 alkaline. Derivation Chain: Why 0.1 Units Matters pH is a log of $[\text{H}^+]$ $\rightarrow$ each whole unit = 10$\times$ change in $[\text{H}^+]$ $\rightarrow$ 0.1 unit $\approx$ 26% change in $[\text{H}^+]$ $\rightarrow$ normal 7.35–7.45 is a genuinely tight band in $[\text{H}^+]$ terms Common Errors • Error: Reading pH linearly, treating 7.0 and 7.4 as nearly the same. They differ ~ 2.5-fold in $[\text{H}^+]$. • Error: Assuming that neutral water at pH 7.0 has no protons. At pH 7.0 the $[\text{H}^+] = 10^{-7}\text{M}$. • Error: Assuming an alkaline solution has no protons. Even at pH 11.5 protons are present, just very few. Prediction Prompt • Predict: Arterial pH falls from 7.40 to 7.30. Predict roughly how much $[\text{H}^+]$ has changed and whether or not this is clinically trivial.

Section 4: Metabolism Is the Acid Source

Instructor Narrative

The move: Except for external intervention, every proton the lungs and kidneys clear was produced by metabolism, and the nutrient fuel a cell oxidizes predicts the acid it produces. Establishing the source first makes the rest of the chapter the removal of that acid.

Each fuel makes a characteristic acid

Carbohydrate catabolism yields carbonic acid aerobically (from CO_2 hydration) and lactic acid anaerobically. Fatty acid oxidation produces the ketoacids acetoacetic and β -hydroxybutyric acid. Amino acid catabolism produces ammonium (NH_4^+) and α -keto acids, and the hydrolysis of nucleotides and sulfur-containing amino acids releases phosphoric and sulfuric acid. This list is not for memorization; the *pattern* is the point: pathway predicts product.

Cells are the body's main proton factory

Roughly forty trillion cells, whose collective intracellular fluid is about forty percent of body weight, are all running these reactions, which is why the chapter treats cells as the origin of bodily H^+ and tackles intracellular regulation first. A clinically important case to flag now: in untreated type 1 diabetes, cells that cannot take up glucose shift to fatty acid oxidation, accumulate ketoacids, and develop a metabolic acidosis, the acid supply outpacing removal.

Volatile vs. fixed acid sets up the two organs

This chapter focuses on acid products that can be sorted into the two categories. First, there is carbonic acid which forms from the hydration of CO_2 . This acid is volatile: depending on the direction its reaction runs, it can become CO_2 and leave through the lungs or it can dissociate into protons and HCO_3^- (which can be selectively retained or excreted by the kidneys as needed). The other metabolic acids (lactic, keto, phosphoric, sulfuric) are fixed in that they cannot be exhaled at the lungs and can only be removed via the urine.

Structured Outline / Causal Chain

Pathway Predicts the Acid

- Aerobic carbohydrate $\rightarrow$ carbonic acid (from CO_2); anaerobic $\rightarrow$ lactic acid.
- Fatty acid oxidation $\rightarrow$ acetoacetic and β -hydroxybutyric acid (ketoacids).
- Amino acid catabolism $\rightarrow$ NH_4^+ and α -keto acids; nucleotide / sulfur-containing amino acid hydrolysis $\rightarrow$ phosphoric, sulfuric acid.

Volatile vs. Fixed (the setup for two organs)

- Volatile: carbonic acid $\rightarrow$ CO_2 , removable by the lungs.
- Fixed: lactic, keto, phosphoric, sulfuric; only the kidney removes these.

Common Errors

- **Error:** Treating all metabolic acids as substances the lungs can handle. Only the volatile (carbonic) fraction leaves through breathing.
- **Error:** Forgetting cells are the acid source, then being surprised that consideration of ICF regulation comes first.

Prediction Prompt

- **Predict:** A cell switches from glucose to heavy fatty-acid oxidation. Predict the acids produced and the likely whole-body acid-base shift.

Section 5: Chemical Buffers — The Shock Absorbers That Hide the Problem

First lines of defense. Plant the shock absorber metaphor early: Buffers act instantaneously, but conceal protons rather than removing them

Instructor Narrative	Structured Outline / Causal Chain
The move: Buffers blunt pH changes without removing protons from the body. These shock absorbers soften the bump rather than filling the hole in the road, and the image is worth returning to throughout the section. Later errors about compensation trace back to forgetting that buffering hides protons without eliminating them. A buffer is a weak-acid / conjugate-base pair that trades protons A buffer is a pair, a weak acid and its conjugate base, that binds H^+ when protons rise and releases H^+ when they fall. Intracellularly, the two big buffer systems are phosphate (Na_2HPO_4 / NaH_2PO_4) and protein (where the work is done mostly by the imidazole group of histidine flipping between protonated and unprotonated near physiological pH). The reactions run by the law of mass action: add protons and the equilibrium shifts to bind them; remove protons and there's a shift to release them. Hiding, not fixing, and almost instantaneous The crucial limit: buffering is extraordinarily fast and can bind the large majority of added protons, but it never removes them from the body. A buffered proton is hidden, not gone, and it remains in the body until the lungs or kidneys remove it. Buffers act like shock absorbers: they minimize the pH change the body undergoes, but the acid load is still present. Buffer systems act together, and two non-chemical mechanisms add capacity The isohydric principle says every buffer system present shares the same H^+s, so they all shift together and a challenge is spread across all of	What a Buffer Is and Does • A weak acid plus its conjugate base; binds H^+ when high, releases when low. • ICF systems: phosphate (Na_2HPO_4 / NaH_2PO_4) and protein (histidine imidazole). • Acts almost instantly; can bind most added protons. The Shock-Absorber Limit • Buffers hide protons; they do not remove them from the body. • A buffered proton must still be excreted by the kidney (or exhaled as CO_2). Isohydric Principle + Non-Chemical Buffering • All buffer systems share the same H^+ and shift together, so a load is distributed across all of them. • Measure one pair → infer overall acid-base status (isohydric principle). • Metabolic buffering: adjust acid production via enzymes. Organellar: sequester H^+ in organelles. Common Errors • Error: Believing buffering corrects pH. It only blunts the change; the protons are still in the body. • Error: Thinking one buffer system dominates. The isohydric principle means they act in concert. Prediction Prompt

them. This principle is also clinically useful: measure one accessible pair (like CO_2 and HCO_3^-) and you have effectively read the status of all of them. Two non-chemical mechanisms add capacity: metabolic buffering (dialing acid production up or down via enzyme activity) and organellar buffering (sequestering protons inside lysosomes or mitochondria, away from cytoplasmic proteins).

- **Predict:** ICF pH drops sharply. Predict the direction of the phosphate reaction and which protein residue does most of the binding.

Section 6: Acid Extruders, Acid Loaders, and the ICF–ECF Handoff

Instructor Narrative

The move: When buffering is no longer sufficient, which is nearly always the case, the cell moves protons across its membrane by transport, which necessarily shifts the acid load into the ECF. This is where the analysis moves from a single cell to the whole body.

Transporters move protons both ways

When buffering is not enough, cells transport protons across the cell membrane. Acid extruders push H^+ out (ATP-driven proton pumps, or Na^+ -coupled exchangers riding the sodium gradient) and pull base anions in; functionally, importing HCO_3^- counts as extruding acid because the base will bind protons. Acid loaders run the reverse when ICF pH climbs too high. Many of these transporters move equal charge in both directions so they stay electroneutral and do not disturb the membrane voltage.

The cell is passively acidified, so active extrusion is required

The direction of net proton movement is counterintuitive. ICF pH (≈ 6.8 – 7.1) is more acidic than that of the ECF (≈ 7.4), so H^+ concentration is higher inside: one would expect protons to leak out. But the negative membrane voltage pulls positive protons inward, and that electrical driving force is the larger of the two: a membrane potential near -70 mV outweighs the chemical gradient favoring H^+ efflux, so net proton movement is inward. HCO_3^- simultaneously leaks out down its gradient.

Structured Outline / Causal Chain

Two Directions of Transport

- Acid extruders: H^+ out (proton pumps; Na^+ -driven exchange) and base anions in; ICF pH rises.
- Importing HCO_3^- = functional acid extrusion (because the base binds H^+).
- Acid loaders: the reverse, used when ICF pH is too high. pH falls (these are often electroneutral exchanges).

Causal Chain: Why the Cell Is Passively Acidified

ICF $[\text{H}^+] > \text{ECF } [\text{H}^+]$, so the chemical gradient favors H^+ exit, while the inside-negative membrane voltage favors H^+ entry, and the electrical force is the larger of the two, so net H^+ movement is inward, and HCO_3^- simultaneously leaks outward down its own gradient, removing base $\rightarrow$ ICF pH would fall continuously. Active extrusion of H^+ is therefore required to hold it.

ECF as Middleman

- ICF problems are handed to the ECF (interstitium = medium; blood = transport to exchange organs/environment).

The net result is continuous passive acidification of the cell, layered on top of its own metabolic acid production. That is why active extrusion and buffering are not optional, they are essential.

ECF is the middleman

Cells transfer their acid-base load into the ECF, which becomes both the medium (interstitial fluid bathing the cells) and the transport route (blood) to the lungs and kidneys. The reverse is clinically relevant: if *those* organs malfunction and ECF pH drifts, the cells can no longer export protons correctly and ICF protein function suffers. The two compartments are coupled.

- Organ dysfunction → ECF pH falls → the outward driving force for H^+ extrusion falls → ICF cannot unload → ICF pH falls → protein dysfunction.

Common Errors

- **Error:** Predicting net proton efflux from the concentration gradient alone. In this instance, membrane voltage actually reverses the net direction.
- **Error:** Treating ICF and ECF pH as independent. They are continuously coupled.

Prediction Prompt

- **Predict:** A drug blocks the Na^+-H^+ exchanger. Predict the effect on ICF pH and on the cell's reliance on chemical buffering.

Section 7: The Bicarbonate System — One Equation, Two Organs

The chapter's central spine. The carbonic acid equation is the one relationship every later section points back to.

Instructor Narrative	Structured Outline / Causal Chain
The move: <i>The carbonic acid equation is the central relationship in the chapter, because it is the only buffer system whose two ends are controlled by two different organs. The lungs control the CO_2 end; the kidneys control the HCO_3^- end.</i> The equation and why carbonic anhydrase is so important The dominant ECF buffer is the bicarbonate system: $CO_2 + H_2O \leftrightarrow H_2CO_3 \leftrightarrow H^+ + HCO_3^-$. It carries roughly 80% of ECF buffering capacity. The reaction happens everywhere but is enormously accelerated by carbonic anhydrase at the two sites where it matters most: in red blood cells, which load CO_2 for transport and unload it for exhalation at the lungs, and in renal tubule cells, which set the ratio of H^+ to HCO_3^-	The Carbonic Acid Equation • $CO_2 + H_2O \leftrightarrow H_2CO_3 \leftrightarrow H^+ + HCO_3^-$. • ~ 80% of ECF buffering; accelerated by carbonic anhydrase in RBCs and renal cells. • H_2CO_3 is the pivot: forms during acid buffering, dissociates during base buffering. Causal Chain: Strong to Weak Conversion (acid challenge) Strong HCl added → H^+ bound by HCO_3^- → forms weak acid H_2CO_3 (dissociates much less) → pH falls only slightly instead of crashing Two Organs, Two Ends

excretion. Carbonic acid sits in the middle of the system as the pivot: it forms when acid is buffered and dissociates when base is buffered.

Strong acids and bases are converted into weak ones

Take the two challenges in turn. Infuse strong HCl and the protons are bound by HCO_3^- to form H_2CO_3 : a strong, fully dissociating acid has become a weak, barely dissociating one, so pH barely moves. Add strong NaOH and the OH^- removes a proton from H_2CO_3 to make water plus the weak base NaHCO_3 . Either way the system trades a large pH change for a small one, exactly the weak-acid logic discussed earlier.

Two adjustable ends is what makes it powerful

The reason this buffer outperforms others is not its chemistry but its regulation. The pK_a for the bicarbonate system (6.1 at human body temperature) represents the pH at which the system is best “poised” to mediate either an acidic or basic challenge. There are equal concentrations of HCO_3^- and CO_2 at that point (ratio = 1, the log of 1 = 0, therefore, $\text{pH} = \text{pK}_a + 0$). Because of this, the system is better adapted to work at pHs much less than the normal 7.40. Keep in mind, though, that both ends of the bicarbonate system are *adjustable*, and that’s what makes this system so powerful. The lungs control the left side by regulating how much CO_2 leaves the body, and the kidneys control the right side by regulating H^+ and HCO_3^- excretion and production. A buffer with two independently controlled ends can be steered, which is why the rest of the chapter is about which organ adjusts which end.

- Lungs control the left side: CO_2 removal.
- Kidneys control the right side: H^+ and HCO_3^- handling.
- Independently adjustable ends → this buffer can be actively steered as appropriate.

Common Errors

- **Error:** Thinking bicarbonate is effective because it is the strongest buffer. Its pK_a (6.1) is far from pH 7.4: This buffer is effective because both ends are regulated.
- **Error:** Forgetting carbonic anhydrase. Without it the CO_2 hydration reaction is too slow to matter at RBCs and tubule cells.

Prediction Prompt

- **Predict:** Blood CO_2 rises. Using the equation, predict the direction of the shift and what happens to H^+ and HCO_3^- .

Section 8: The Henderson-Hasselbalch Equation — Reading and Predicting Compensation

Instructor Narrative

The move: The Henderson-Hasselbalch equation is used here to predict pulmonary and/or renal compensation. The one substitution worth explaining is the replacement of H_2CO_3 , which is difficult to measure, with

Structured Outline / Causal Chain

The Equation and Its One Substitution

- $K_a = ([\text{H}^+][\text{HCO}_3^-])/[\text{H}_2\text{CO}_3]$; H_2CO_3 hard to measure.

dissolved CO_2 , which is calculated from PCO_2 . After that, the equation gives the variable each organ moves and the direction it needs to move it.

Why we write the equation in terms of CO_2 instead of H_2CO_3

Start from $K_a = ([\text{H}^+][\text{HCO}_3^-])/[\text{H}_2\text{CO}_3]$. H_2CO_3 is essentially impossible to measure, whereas dissolved CO_2 is readily calculated from blood PCO_2 times a solubility coefficient. Since hydrated and free CO_2 sit in a fixed 1:1000 ratio, one can substitute CO_2 for H_2CO_3 (rolling the factor into a modified constant) and rearrange to the form clinicians use: $\text{pH} = \text{pK}_a + \log([\text{HCO}_3^-]/[\text{CO}_2])$, with $\text{pK}_a = 6.1$ in blood.

Two knowns give the third variable, and predict compensation

The practical power is twofold. Given any two of pH , HCO_3^- , and CO_2 , one can easily solve for the third, which is exactly what an arterial blood gas exploits. And the ratio form predicts the response to a disturbance: in acidosis the body raises HCO_3^- (kidney) and lowers CO_2 (lungs); in alkalosis it does the reverse. The equation names which variable each organ moves, and in which direction.

Poised low, but steerable everywhere

The system buffers best where HCO_3^- and CO_2 are equal, at $\text{pH} = \text{pK}_a = 6.1$, well below the body's 7.4, so by classic buffer logic bicarbonate appears poorly suited. The resolution is the same as before: because both ends are adjustable, the body is not stuck at the poised point. It holds pH at 7.4 by keeping the HCO_3^- -to- CO_2 ratio near 20:1 and shifting it on demand.

- Hydrated : free $\text{CO}_2 \approx 1:1000 \rightarrow$ substitute CO_2 for H_2CO_3 (modified constant).
- $\text{pH} = \text{pK}_a + \log([\text{HCO}_3^-]/[\text{CO}_2])$; $\text{pK}_a = 6.1$; $[\text{CO}_2] = \text{solubility} \times \text{PCO}_2$.

Causal Chain: From Disturbance to Compensation

Acidosis (low pH) $\rightarrow$ both ends are adjusted in parallel: the lungs lower $[\text{CO}_2]$ within seconds (increase ventilation) and the kidney raises $[\text{HCO}_3^-]$ over hours (reduce excretion, make new). The HCO_3^- -to- CO_2 ratio rises, and by Henderson-Hasselbalch that is a pH returning toward 7.4.

The Poised-Point Paradox

- Best buffering at $\text{pH} = \text{pK}_a = 6.1$ (ratio = 1), far from 7.4.
- But both ends are adjustable, so the body steers the ratio ($\sim 20:1$ at pH 7.4).

Common Errors

- **Error:** Thinking of bicarbonate as a poor buffer because pK_a is 6.1 (far removed from the normal blood pH of ~ 7.4). Adjustability, not proximity to 7.4, is the point.
- **Error:** Forgetting $[\text{CO}_2]$ must come from $\text{PCO}_2 \times \text{solubility}$ before using the equation.

Prediction Prompt

- **Predict:** An ABG gives pH and PCO_2 only. Predict how you would obtain $[\text{HCO}_3^-]$ and what a low value alongside low pH suggests.

Section 9: The Fast Defense — Respiratory Control of CO_2

Second line of defense (seconds). The lungs steer the left side of the bicarbonate equation by changing how much CO_2 stays in the blood.

Instructor Narrative

The move: Breathing is connected to pH through chemoreceptors. The lungs do not sense pH themselves; chemoreceptors detect H^+ and CO_2 , and the brainstem adjusts ventilation. The same loop, driven by something other than blood chemistry, can produce the pH disturbance instead of correcting it.

Hypercapnia lowers pH; chemoreceptors detect it

When metabolism increases, CO_2 production climbs, PCO_2 rises (hypercapnia), the equation shifts right, and pH falls. Two sensor populations report this. Central chemoreceptors in the ventral medulla actually sense H^+ , but as a proxy for CO_2 : blood CO_2 , being lipophilic, crosses the blood-brain barrier, drives the carbonic acid reaction in the brain's interstitium and CSF, and frees protons. Because CSF has almost no protein and thus little buffering, those protons stay free and the receptors are exquisitely CO_2 -sensitive. Peripheral chemoreceptors, glomus cells in the carotid and aortic bodies, sense low O_2 , high CO_2 , and low pH, with hypoxemia their strongest trigger.

The brainstem adjusts ventilation

Receptor signals feed the medullary respiratory center (the dorsal and ventral respiratory groups), which drives the diaphragm to change the rate and depth of breathing. Acidosis stimulates hyperventilation, which blows off CO_2 , shifts the equation left, and increases pH back toward normal within seconds. Alkalosis does the opposite via hypoventilation. Pulmonary compensation is fast but rarely perfectly complete.

When the lung is the problem, not the fix

The loop also runs in reverse. If ventilation is driven by something other than blood chemistry (anxiety, fever, altitude, or lung disease), it can itself create the disturbance. Chronic hypoventilation traps CO_2 and causes respiratory acidosis; chronic hyperventilation washes out CO_2 and causes respiratory alkalosis. The same machinery is now the *cause* rather than the correction.

Structured Outline / Causal Chain

Two Chemoreceptor Populations

- Central (ventral medulla): sense H^+ as a proxy for CO_2 ; CO_2 crosses the BBB, CSF has low buffering $\rightarrow$ very CO_2 -sensitive.
- Peripheral (carotid + aortic glomus cells): sense low O_2 , high CO_2 , low pH; hypoxemia is the strongest stimulus.

Causal Chain: The Respiratory Compensation Loop

$\uparrow CO_2$ (hypercapnia) $\rightarrow$ $\downarrow$ pH $\rightarrow$ chemoreceptors detect the change $\rightarrow$ medullary respiratory center $\uparrow$ rate and depth $\rightarrow$ $\uparrow$ ventilation blows off CO_2 $\rightarrow$ equation shifts left $\rightarrow$ $[H^+]$ falls, which is a pH rising toward normal

Same Loop, Run Backward = Disease

- Chronic hypoventilation $\rightarrow$ CO_2 retention $\rightarrow$ respiratory acidosis.
- Chronic hyperventilation $\rightarrow$ CO_2 washout $\rightarrow$ respiratory alkalosis.

Common Errors

- **Error:** Believing central receptors sense CO_2 directly. They sense H^+ ; CO_2 is the upstream driver via the BBB.
- **Error:** Assuming ventilation only *compensates* acidosis or alkalosis. These disorders can be *caused* by respiratory disturbances.

Prediction Prompt

- **Predict:** A panicked student hyperventilates before an exam. Predict the direction of change in PCO_2 and arterial pH, and name the disturbance.

Section 10: The Potent Defense — Why Only the Kidneys Eliminate Fixed Acid

Third line of defense (hours). Slow, but the only system that truly removes fixed acid and rebuilds buffer.

Instructor Narrative	Structured Outline / Causal Chain
The move: <i>The division of labor is explicit: buffers hide protons, the lungs exhale volatile “acid”, and only the kidney removes fixed acid from the body and synthesizes new buffer. The kidney is the slowest of the three defenses, but the only one that resolves a fixed-acid load completely.</i> Only the kidney eliminates acid rather than just hiding it The hierarchy runs in a timeline: fast and local (buffers), then faster systemic (lungs in seconds), then slow systemic (kidneys in minutes to hours). The kidney is last and slowest but most potent because it is the only step that actually takes protons out of the body, in acidic urine, and the only one that can build new buffer. Buffered protons were only temporarily bound to buffers and CO_2 contains no dissociable protons; the kidney is where they are finally removed. Filtration sets the stage; reabsorption and secretion do the work The clearance logic applies here: at a given GFR a fixed fraction of a filterable solute enters the tubular fluid, but what is actually removed depends on reabsorption and secretion. HCO_3^- is freely filtered, about 4,320 mmol per day, and almost none should appear in urine, so the nephron must reabsorb nearly all of it continuously. H^+s, by contrast, are largely buffered by plasma proteins and escape filtration, so the kidney must secrete them. The power of renal acid-base control is the ability to tune three events at once: secrete H^+, reabsorb filtered HCO_3^-, and make new HCO_3^-. New bicarbonate, not reabsorption, restores consumed buffer Reabsorbing filtered HCO_3^- only prevents loss; it cannot replace buffer already consumed neutralizing fixed acid. To keep the ECF buffer pool	The Defense Hierarchy (timeline) • Buffers: instant, local, hide protons. • Lungs: seconds, blow off volatile (CO_2) acid. • Kidneys: minutes–hours, eliminate fixed acid in urine and build new buffer. Clearance Logic for Acid-Base • HCO_3^-: freely filtered ($\sim 4,320$ mmol/day), nearly all reabsorbed. • H^+s: buffered by plasma proteins, escape filtration → must be secreted. • Three tunable actions: secrete H^+, reabsorb HCO_3^-, make new HCO_3^-. Common Errors • Error: Thinking the lungs can eliminate a fixed-acid load. They only remove the volatile CO_2 fraction. • Error: Treating HCO_3^- reabsorption and new HCO_3^- production as the same thing. One prevents loss; the other restocks buffer. Prediction Prompts • Predict: A patient has a chronic fixed-acid load. Predict what happens to arterial pH, PCO_2, and the ECF bicarbonate pool over the first minutes, and then over the following hours. Name which defense is responsible at each stage. • Predict: A second patient with a chronic fixed-acid load presents with an arterial pH of 7.38. Explain why

from depleting over days of net acid production, the kidney must synthesize new HCO_3^- and add it to the blood. Reabsorption defends the existing supply; new production restocks it.

this patient may still be in serious trouble, and what single measurement would reveal it.

Section 11: The Proximal Tubule — Reabsorbing Bicarbonate by Secreting Protons

Instructor Narrative

The move: Most of the protons the proximal tubule secretes do not acidify the urine; they help recover filtered bicarbonate. The route is indirect because filtered HCO_3^- cannot cross the apical membrane and must be converted to CO_2 and then rebuilt inside the cell.

Secretion is indirect, and it starts with CO_2

Protons are not transported directly across the PCT wall. Inside tubular cells, CO_2 (from peritubular blood, from cell metabolism, or from the tubular fluid) is rapidly hydrated via carbonic anhydrase to H_2CO_3 , which dissociates to H^+ and HCO_3^- . The basolateral $\text{Na}^+\text{-K}^+$ pump builds the sodium gradient that powers an apical $\text{Na}^+\text{-H}^+$ exchanger, driving protons into the lumen, with extra protons pushed by an ATP-dependent pump. The HCO_3^- made in the same step is reabsorbed toward blood, in part by cotransport with Na^+ and counter-transport with Cl^- .

Why the proton has to go first

Filtered HCO_3^- cannot cross the apical membrane directly. So, a secreted proton meets filtered HCO_3^- in the lumen, forms H_2CO_3 , which becomes CO_2 and water; that CO_2 diffuses into the cell, where carbonic anhydrase regenerates H^+ and HCO_3^- , and the HCO_3^- is reabsorbed. The proton is then recycled to bind the next bicarbonate. That means proton secretion's major function is to facilitate bicarbonate reabsorption; acid excretion is the minor side effect, and PCT H^+ secretion and HCO_3^- reabsorption run about 1:1.

Structured Outline / Causal Chain

Secretion Serves Reabsorption

- Most PCT H^+ secretion serves mainly HCO_3^- reabsorption, not urine acidification.
- Filtered HCO_3^- cannot cross the membrane directly: it is converted, then regenerated.
- H^+ secretion and HCO_3^- reabsorption run ~ 1:1 in the PCT.

Causal Chain: How Filtered Bicarbonate Is Reabsorbed

H^+ secreted into lumen ($\text{Na}^+\text{-H}^+$ exchange, powered by $\text{Na}^+\text{-K}^+$ pump) $\rightarrow \text{H}^+ + \text{filtered HCO}_3^- \rightarrow \text{H}_2\text{CO}_3 \rightarrow \text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{CO}_2$ diffuses into the cell $\rightarrow$ carbonic anhydrase regenerates $\text{H}^+ + \text{HCO}_3^- \rightarrow \text{HCO}_3^-$ reabsorbed to blood; H^+ recycled to lumen

New Bicarbonate

- When the filtered, then reabsorbed, HCO_3^- supply is exhausted, blood CO_2 feeds continued intracellular HCO_3^- production.
- This HCO_3^- is new ECF buffer, not recovered filtrate.

Common Errors

- **Error:** Thinking filtered bicarbonate is reabsorbed intact. It combines with H^+ in the tubular lumen to form H_2CO_3 which separates into CO_2 and H_2O . The CO_2 then diffuses into the cell and is rehydrated to form

Making new bicarbonate when filtered, then reabsorbed, buffer runs out

During a metabolic acid load, all filtered HCO_3^- can be reabsorbed and the kidney still needs to keep secreting protons. As long as blood supplies CO_2 , the intracellular reaction keeps regenerating HCO_3^- that is genuinely new, added to the ECF rather than merely recovered. That is the production step from the previous section, stated as a mechanism.

H_2CO_3 which dissociates into HCO_3^- and H^+ . The HCO_3^- is absorbed into the blood and the H^+ is resecreted.

- **Error:** Assuming proton secretion mainly acidifies urine. In the PCT its main effect is bicarbonate recovery.

Prediction Prompt

Predict: A carbonic anhydrase inhibitor is given. Predict the effect on HCO_3^- reabsorption, urine pH, and ECF pH.

Section 12: The Late Nephron — Intercalated Cells and Urine Buffers

Distal tubule fine-tuning layered on proximal tubule bulk recovery.

Instructor Narrative	Structured Outline / Causal Chain
The move: The two intercalated cell types run the same carbonic acid reaction in opposite directions, sending products opposite ways. The urine also requires its own non-bicarbonate buffers, because without them the falling luminal pH stalls further proton secretion. Type A and type B are mirror images Late fine-tuning occurs at intercalated cells scattered among principal cells in the late DCT and early collecting duct. Both run the carbonic acid reaction; they differ only in what they do with the products. Type A cells handle acidosis: apical ATP-dependent pumps secrete H^+ into the lumen while HCO_3^- goes to blood, raising ECF pH. Type B cells handle alkalosis: they are inverted type A cells, secreting HCO_3^- into the lumen and returning H^+ to blood, lowering ECF pH. Diet shows up here: a vegetable-rich, low-meat diet drives type B activity and alkaline urine. Why urine needs phosphate and ammonia buffers There is a physical limit. As urine acidifies (normally near pH 6, occasionally as low as ~ 4.6), the gradient opposing further proton	Intercalated Cell Subtypes • Both run $\text{CO}_2 + \text{H}_2\text{O} \leftrightarrow \text{H}_2\text{CO}_3 \leftrightarrow \text{H}^+ + \text{HCO}_3^-$; they differ only in details of product handling and shipping. • Type A (acidosis): secrete H^+ to lumen, HCO_3^- to blood; urine pH can fall to ~ 4.6. • Type B (alkalosis): secrete HCO_3^- to lumen, H^+ to blood; urine pH can rise to ~ 8.0. Why Urine Needs Buffers • Decreasing urine pH steepens the gradient opposing H^+ secretion (stalls transporters). • $\text{HPO}_4^{2-} + \text{H}^+ \rightarrow \text{H}_2\text{PO}_4^-$; $\text{NH}_3 + \text{H}^+ \rightarrow \text{NH}_4^+$ trap protons in urine. • Buffered secretion lets the cell keep secreting and regenerating HCO_3^-. Causal Chain: Glutamine: Two Products

secretion steepens and the transporters stall. Tubular buffers solve this: $\text{HPO}_4^{2-} + \text{H}^+ \rightarrow \text{H}_2\text{PO}_4^-$ and $\text{NH}_3 + \text{H}^+ \rightarrow \text{NH}_4^+$. A secreted proton bound to a non-bicarbonate buffer stays in the urine to be excreted, keeps luminal pH from crashing, and lets secretion (and HCO_3^- regeneration) continue. A large fraction of acid excretion leaves as phosphate- and ammonia-buffered protons.

Glutamine: buffer and new bicarbonate at once

The kidney metabolizes the amino acid glutamine to make NH_4^+ (which carries protons out in the urine) and HCO_3^- , which is absorbed into blood as yet another source of new buffer. One reaction both excretes acid and restocks the ECF buffer pool.

Kidney metabolizes glutamine $\rightarrow$ produces NH_4^+ and $\text{HCO}_3^- \rightarrow \text{NH}_4^+$ carries a proton into the urine and is excreted; HCO_3^- is absorbed to blood. That is acid excreted and new buffer made in one step.

Common Errors

- **Error:** Treating type A and B cells as unrelated mechanisms. Type B is an inverted type A; mass action sets the direction.
- **Error:** Forgetting urine buffers. Without phosphate and ammonia, urine pH would crash and stop secretion.

Prediction Prompt

- **Predict:** Urinary phosphate and ammonia fall. Predict the effect on urine pH and on the proton gradient across the tubule wall.

Section 13: Naming the Disturbance — Respiratory vs. Metabolic, Acidosis vs. Alkalosis

Instructor Narrative	Structured Outline / Causal Chain
The move: Two questions fill the 2×2 classification: pH gives the direction, and whether PCO_2 or HCO_3^- accounts for it gives the cause. Respiratory names the lung-controlled end of the equation; metabolic names other pH disturbances. Two axes, four boxes Arterial pH below 7.35 is acidosis; above 7.45 is alkalosis: that is the first axis. The second axis is cause. If the abnormality tracks PCO_2 (the lung-controlled end), it is respiratory; if it tracks HCO_3^- (everything else), it is metabolic. So, acidosis with high PCO_2 (above ~ 40 mmHg) is respiratory acidosis, the lungs failing to clear CO_2; acidosis with low HCO_3^- (below ~ 24 mEq/L) is metabolic acidosis. Alkalosis splits the same way: low PCO_2 is respiratory, high HCO_3^- is metabolic.	The 2×2 Decision Rule • (1 or 2, directions); pH < 7.35 = acidosis; > 7.45 = alkalosis. • (3 or 4, causes); Tracks $\text{PCO}_2 \rightarrow$ respiratory; tracks $\text{HCO}_3^- \rightarrow$ metabolic. • Respiratory acidosis = $\uparrow\text{PCO}_2$; metabolic acidosis = $\downarrow\text{HCO}_3^-$ (alkalosis mirrors both). Causal Chain: Why Metabolic Acidosis Shows Low HCO_3^- Fixed acid adds $\text{H}^+ \rightarrow \text{HCO}_3^-$ buffers it and is consumed, so measured $[\text{HCO}_3^-]$ falls. A low HCO_3^- is the fingerprint of acid buffering. The General Equivalence

Metabolic acidosis from two directions

Metabolic acidosis can appear as either high H^+ or low HCO_3^- , and both are the same event seen from two sides. Added protons are buffered by HCO_3^- , consuming them, so measured HCO_3^- falls: a low value is the fingerprint of buffering against an acid load. The general rule: with respect to pH, losing base is equivalent to gaining acid, and vice versa.

- For pH, loss of base = gain of acid (and vice versa).
- High H^+ and low HCO_3^- describe the same metabolic acidosis.

Common Errors

- **Error:** Confusing the direction (pH) with the cause (PCO_2 vs HCO_3^-). One needs both to name the disorder.
- **Error:** Thinking high H^+ and low HCO_3^- are two different problems. They are one event from two sides.

Prediction Prompt

- **Predict:** A patient has pH 7.30 and PCO_2 55 mmHg. Predict the disorder and which organ is implicated.

Section 14: Reading the Blood — Anion Gap, ABGs, and the Davenport Diagram

Instructor Narrative

***The move:** Three tools read the blood: anion gap sub-classifies metabolic acidosis, arterial blood gas supplies pH and PCO_2 , and the Davenport diagram displays compensation as movement on a graph. Each disorder is a readout of an underlying metabolic, pulmonary, or renal problem rather than a disease in itself.*

The anion gap exploits electroneutrality

Real solutions are electroneutral, total cations equal total anions, so a true gap is zero. The clinical anion gap, $AG = ([Na^+] + [K^+]) - ([Cl^-] + [HCO_3^-])$, counts only the major ions and lands near 15 mM, the “gap” being unmeasured anions. Its use: when excess metabolic acid is buffered, HCO_3^- falls without Cl^- rising, so the gap widens, a high-anion-gap metabolic acidosis (lactic acidosis, ketoacidosis). If instead HCO_3^- is lost and Cl^- replaces it (GI or renal HCO_3^- loss), the gap stays normal, a non-anion-gap acidosis. A single electrolyte panel separates the two.

Structured Outline / Causal Chain

Anion Gap

- Electroneutrality: the true cation-anion gap is zero; AG counts only major ions.
- $AG = ([Na^+] + [K^+]) - ([Cl^-] + [HCO_3^-]) \approx 15$ mM normally.
- High-AG acidosis: acid buffered, $HCO_3^- \downarrow$ without $Cl^- \uparrow$ (lactic, keto). Non-AG acidosis: HCO_3^- lost, Cl^- replaces it (GI / renal).

Arterial Blood Gas

- Measures pH, PCO_2 , PO_2 ; venous gas (VBG) parallels the trend.
- $pH + PCO_2 \rightarrow [HCO_3^-]$ via Henderson-Hasselbalch.

The ABG gives the numbers

Arterial blood gas analysis (drawn from an artery, technically more difficult and painful than the venous sampling sometimes substituted) measures pH, PCO_2 , and PO_2 . With pH and PCO_2 in hand, Henderson-Hasselbalch yields HCO_3^- , and the three together let one read respiratory function, pH status, and the likely source of a disturbance.

The Davenport diagram makes compensation visible

Davenport's plot puts HCO_3^- on the y-axis and pH on the x-axis, with CO_2 isopleths (curved PCO_2 lines) and a buffer line whose slope reflects blood buffering capacity, mostly hemoglobin, tying back to how blood carries CO_2 . Two rules to read it: at a fixed PCO_2 you scan along one curve; changing buffering moves you onto a different buffer line. A pure rise in PCO_2 from say 40 mmHg to 60 mmHg shifts the assessment toward respiratory acidosis (with HCO_3^- nudging up, as the equation predicts); falling PCO_2 moves toward respiratory alkalosis; vertical steps at a fixed PCO_2 are metabolic. Chronic compensation appears as a second arrow returning pH toward 7.4, renal for respiratory problems, respiratory for metabolic ones.

Disorders are symptoms, simple or mixed

Acid-base disorders are not diseases in themselves; they are readouts of metabolic, pulmonary, or renal dysfunction (or of drugs, altitude, anxiety). A single cause is a simple disorder; overlapping causes, say respiratory plus renal failure, make a mixed disorder that the Davenport diagram can reveal, but only with a seasoned user's experience. The chapter's payoff is being able to look at three numbers and reconstruct which of the body's three defenses is failing and which is compensating.

Causal Chain: Why Acute Respiratory Acidosis Moves Along the Buffer Line

PCO_2 rises from 40 to 60 mmHg → the carbonic acid equation is driven right → both H^+ and HCO_3^- are generated, so pH falls while HCO_3^- rises slightly. On the diagram that is the point moving left and slightly up along the buffer line, landing on the 60 mmHg isopleth → over hours the kidney secretes H^+ and synthesizes new HCO_3^- → HCO_3^- rises above the original buffer line and pH returns toward 7.4

Reading the Davenport Diagram

Rules to follow: at a fixed PCO_2 you read along one curve; changing buffering moves you onto a different buffer line. A pure rise in PCO_2 slides toward respiratory acidosis, falling PCO_2 moves toward respiratory alkalosis. Vertical moves at a fixed PCO_2 are metabolic. Chronic compensation appears as a second arrow returning pH toward 7.4

Disorders Are Symptoms

- Not diseases per se, readouts of metabolic, pulmonary, or renal dysfunction (or drugs / altitude / anxiety).
- Simple = one cause; mixed = overlapping causes (e.g., respiratory + renal).

Common Errors

- **Error:** Forgetting the buffer line is mostly a function of [hemoglobin]. A different hematocrit changes the line's slope.
- **Error:** Reading a high anion gap as just "acidosis." It specifically flags added-acid (not HCO_3^- loss) metabolic acidosis. Metabolic acidosis can also occur in a non-AG form.

Prediction Prompt

- **Predict:** PCO_2 jumps from 40 mmHg to 60 mmHg acutely. Predict the Davenport move and the direction of the pH change, then the chronic compensating arrow.
- **Predict:** The anion gap is elevated in a patient with acidosis. Predict the likely category of cause and one example.